

Søgeprotokol for Nationale Kliniske Retningslinjer - Guidelines

Projekttitel/aspekt	NKR for Behandling af angst hos børn og unge /
Fagkonsulenter / projektleder	Kristian Bech Arendt / Rasmus Trap Wolf
Søgespecialist	Helene Sognstrup
Senest opdateret	14.01.2016

Baggrund	Forekomsten af angstlidelser for voksne i Danmark er vurderet til at være 13-29 % i livstidsprævalens ¹ . I perioden 2001 til 2011 har der været en stigning i antallet af børn og unge med angsttilstande fra ca. 200 til 450 per år i den regionale psykiatri ² (ambulante og indlagte), og angstlidelser er nogle af de hyppigste tilstande, som ses i børne- og ungdomspsykiatrien....
Søgetermer	Engelske: anxiety, anxiety disorders, anxious, social phobia, social anxiety, generaliz(s)ed anxiety disorder Danske: angst, social fobi, social angst Norske: angst, generalisert angstlidelse, angstlidelse, sosial fobi, sosial angst Svenske: ångeststörning, ångest, social fobi, social ångest, generaliserat ångestsyndrom
Inklusions- og eksklusionskriterier	Sprog: Engelsk, dansk, norsk og svensk År: Sidste 10 år (2005-2015) Publikationstyper: guidelines Population: Børn og unge indtil 18 år

Informationskilder

DATABASER	INTERFACE	FUND	DATO FOR SØGNING
G-I-N International	Internet	34	19.10.2015
NICE (UK)	Internet	5	19.10.2015
National Guideline Clearinghouse (USA)	Internet	22	19.10.2015
Scottish Intercollegiate Guidelines Network (SIGN)	Internet	0	19.10.2015
HTA Databasen (CRD database)	Internet	37	19.10.2015
SBU, Sverige	Internet	5	19.10.2015
Socialstyrelsen, Sverige	Internet	1	19.10.2015
Helsedirektoratet, Norge	Internet	2	19.10.2015
Kunnskapssenteret, Norge	Internet	4	19.10.2015
Netpunkt	Internet	2	19.10.2015

Medline	OVID	72	28.10.2015
Embase	OVID	475	28.10.2015
Cinahl	EBSCO	63	28.10.2015
PsycInfo	OVID	93	28.10.2015

Note:

- Guidelines fra firmaet Hayes er fra valgt for denne NKR
- Søgetermer og inklusions- og eksklusionskriterier er tilpasset de enkelte databaser.
- Doubletetter er så vidt muligt frasorteret ved hjælp af RefWorks. De fundne referencer overføres til fagkonsulenten i Covidence. For nogle af informationskilderne var det ikke muligt at eksportere referencerne til RefWorks; disse er gennemgået af fagkonsulenten i søgeprotokollen og relevante er lagt manuelt ind. De pågældende kilder er markeret med IE (Ikke Eksporteret).
- Fuldtekster præsenteres i Covidence i pdf-format eller som link
- Søgestrategi for hver enkelt database præsenteres – hvis muligt vises det eksplicit hvor mange referencer den enkelte søgestreng genererer
- D. 14-01-2016 er der lavet en supplerende søgning på Separation Anxiety. Der er søgt på Separation anxiety (emneord) OR separation anxiet*. Søgningen gav i alt 6 nye referencer. Se supplerende søgning sidst i dokumentet.

SØGESTRATEGI

Indhold


G-I-N International	3
NICE (UK) (IE)	9
National Guideline Clearinghouse (USA) (IE)	10
Scottish Intercollegiate Guidelines Network (SIGN).....	14
HTA Databasen (CRD database)	14
SBU, Sverige (IE)	19
Socialstyrelsen, Sverige (IE).....	22
Helsedirektoratet, Norge (IE).....	22
Kunnskapssenteret, Norge (IE)	23
Netpunkt.....	25
Medline	26
Embase.....	27
PsycInfo	27
CINAHL	28
Supplerende søgning på Separation Anxiety	29


G-I-N International

(søgt d. 19.10.2015)

Fundne referencer: 34 (Overført til Covidence)

anxiety* OR social phobia OR social anxiety OR anxiety disorder* (dansk, engelsk, norsk, svensk; 2005-2015)


<u>Relevance</u>	<u>Title</u>	<u>Organisation</u>	<u>Type</u>	<u>Date</u>	<u>Relevant Countries</u>	<u>Status</u> 
100%	Social anxiety disorder (CG159)	NICE (UK) - National Institute for Health and Care Excellence	Guideline	May 01, 2013	United Kingdom	Published
65%	Anxiety (CG113)	NICE (UK) - National Institute for Health and Care	Guideline	Jan 01, 2011	United Kingdom	Published


<u>Relevance</u>	<u>Title</u>	<u>Organisation</u>	<u>Type</u>	<u>Date</u>	<u>Relevant Countries</u>	<u>Status</u> 
57%	Anxiety - adults	Excellence SST (DK) - Danish Health and Medicines Authority	Guideline	Nov 28, 2007	Denmark	Published
48%	Anxiety and depression in children and youth— diagnosis and treatment. Medical Services Commission, British Columbia. NGC:009471	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Jan 01, 2010	Canada	Published
48%	Australian clinical practice guidelines: depression and related disorders – anxiety, bipolar disorder and puerperal psychosis – in the perinatal period. A guideline for primary health care professionals. beyondblue: the national depression initiative. NGC:009654	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Feb 01, 2011	Australia	Published
47%	Obsessive compulsive disorder (OCD) and body dysmorphic disorder (BDD) (CG31)	NICE (UK) - National Institute for Health and Care Excellence	Guideline	Nov 01, 2005	United Kingdom	Published
47%	Psychological aspects of chronic pelvic pain. In: Guidelines on chronic pelvic pain. European Association of Urology. NGC:009430	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Feb 01, 2012	International	Published
47%	Screening, Assessment, and Care of Anxiety and Depressive Symptoms in Adults With Cancer: An American Society of Clinical Oncology Guideline Adaptation	ASCO (US) - American Society of Clinical Oncology	Guideline	May 20, 2014	United States	Published
44%	Environmental healthcare design strategies and event	JB1 (AU) - Joanna Briggs	Guideline	Dec 11,	Australia	Published

<u>Relevance</u>	<u>Title</u>	<u>Organisation</u>	<u>Type</u>	<u>Date</u>	<u>Relevant Countries</u>	<u>Status</u> 
44%	related anxiety in pediatric patients (Best Practice 16(11)) Best evidence statement (BEST). Psycho-social interventions for primary caregivers of newly diagnosed pediatric oncology patients. Cincinnati Children's Hospital Medical Center. NGC:009492	Institute AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Nov 28, 2012	United States	Published
41%	Evidence-based practice guideline. Detection and assessment of late life anxiety. University of Iowa Gerontological Nursing Interventions Research Center, Research Translation and Dissemination Core. NGC:006961	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Nov 01, 2008	United States	Published
41%	Putting evidence into practice: evidence-based interventions for anxiety. Oncology Nursing Society. NGC:007728	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Oct 01, 2008	United States	Published
39%	Common mental health disorders (CG123)	NICE (UK) - National Institute for Health and Care Excellence	Guideline	May 01, 2011	United Kingdom	Published
38%	Clinical practice guideline for treatment of patients with anxiety disorders in primary care. GuiaSalud - National Government Agency. Health Technology Assessment Unit, Laín Entralgo Agency. Ministry of Health (Spain). NGC:009479	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Jan 01, 2008	Spain	Published
38%	Post-traumatic stress	NICE (UK) -	Guideline	Mar	United	Published

<u>Relevance</u>	<u>Title</u>	<u>Organisation</u>	<u>Type</u>	<u>Date</u>	<u>Relevant Countries</u>	<u>Status</u> 
	disorder (PTSD) (CG26)	National Institute for Health and Care Excellence		01, 2005	Kingdom	
38%	Practice guideline for the treatment of patients with panic disorder. American Psychiatric Association. NGC:007124	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Jan 01, 2009	United States	Published
37%	Adolescent health risk behavior assessment. Michigan Quality Improvement Consortium. NGC:009445	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Sep 01, 2012	United States	Published
37%	Practice guideline for the treatment of patients with obsessive-compulsive disorder. American Psychiatric Association. NGC:005841	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Jan 01, 2007	United States	Published
34%	Use of psychiatric medications during pregnancy and lactation. American College of Obstetricians and Gynecologists. NGC:006421	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Apr 01, 2008	United States	Published
34%	Practice parameter for psychodynamic psychotherapy with children. American Academy of Child and Adolescent Psychiatry. NGC:009070	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	May 01, 2012	United States	Published
33%	Trauma and post-traumatic stress disorder in patients with HIV/AIDS. New York State Department of Health. NGC:006470	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Dec 01, 2007	United States	Published
32%	Post-traumatic stress disorder. The management of PTSD in adults and children in primary and	AHRQ (US) - Agency for Healthcare Research and	Guideline Clearing Report	Jan 01, 2005	United Kingdom	Published

<u>Relevance</u>	<u>Title</u>	<u>Organisation</u>	<u>Type</u>	<u>Date</u>	<u>Relevant Countries</u>	<u>Status</u> 
	secondary care. National Collaborating Centre for Mental Health. NGC:004204	Quality				
32%	Prevention of secondary disease: mental health care. New York State Department of Health. NGC:005520	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Dec 01, 2006	United States	Published
31%	Practice parameter for the assessment and treatment of children and adolescents with obsessive-compulsive disorder. American Academy of Child and Adolescent Psychiatry. NGC:008982	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Jan 01, 2012	United States	Published
27%	Routine preventive services for children and adolescents (ages 2-21). Michigan Quality Improvement Consortium. NGC:009931	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	May 01, 2013	United States	Published
27%	Clinical Pharmacogenetics Implementation Consortium guideline for CYP2D6 and CYP2C19 genotypes and dosing of tricyclic antidepressants. Clinical Pharmacogenetics Implementation Consortium. NGC:009742	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	May 01, 2013	United States	Published
26%	Practice parameter on gay, lesbian or bisexual sexual orientation, gender nonconformity, and gender discordance in children and adolescents. American Academy of Child and Adolescent Psychiatry. NGC:009316	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Sep 01, 2012	United States	Published
26%	Rehabilitation. In: Clinical guidelines for stroke	AHRQ (US) - Agency for	Guideline Clearing	Sep 01,	Australia	Published

<u>Relevance</u>	<u>Title</u>	<u>Organisation</u>	<u>Type</u>	<u>Date</u>	<u>Relevant Countries</u>	<u>Status</u> 
22%	management 2010. 2005 Sep 8 (revised 2010 Sep). NGC:008108 KSUMC Evidence-Based Clinical Practice Guideline for Management of Bipolar Disorder (KSUMC CPG 1st Edition 2013): Adapted from Canadian Network for Mood and Anxiety Treatments (CANMAT) and International Society for Bipolar Disorders (ISBD) collaborative update of CANMAT guidelines for the management of patients with bipolar disorder: Update 2013 Practice parameter: treatment of nonmotor symptoms of Parkinson disease. Report of the Quality Standards Subcommittee of the American Academy of Neurology. American Academy of Neurology. NGC:007823	Healthcare Research and Quality EBHC KT (SA) - Chair of Evidence-based Healthcare and Knowledge translation, College of Medicine, King Saud University	Report Guideline	2010 Dec 15, 2013	Saudi Arabia	Published
21%	VA/DoD clinical practice guideline for management of post-traumatic stress. Department of Defense. Department of Veterans Affairs. Veterans Health Administration. NGC:008249	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	May 16, 2009	United States	Published
21%	VA/DoD clinical practice guideline for management of post-traumatic stress. Department of Defense. Department of Veterans Affairs. Veterans Health Administration. NGC:008249	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Jan 01, 2010	United States	Published
18%	Music interventions for dental anxiety in paediatric and adult patients (Best Practice 15(2))	JBI (AU) - Joanna Briggs Institute	Guideline	Dec 02, 2011	Australia	Published
15%	Long-term medical management of the pediatric patient after liver	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Aug 01, 2013	United States	Published

<u>Relevance</u>	<u>Title</u>	<u>Organisation</u>	<u>Type</u>	<u>Date</u>	<u>Relevant Countries</u>	<u>Status</u> 
11%	transplantation: 2013 practice guideline by the American Association for the Study of Liver Diseases and the American Society of Transplantation. American Association for the Study of Liver Diseases. NGC:010032 Best evidence statement (BESt). In children with a history of child abuse or neglect does preparation for medical procedures using medical play vs. no preparation reduce anxiety? Cincinnati Children's Hospital Medical Center. NGC:008172	Research and Quality AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	Oct 26, 2010	United Kingdom	Published

NICE (UK) (IE)

(søgt d. 19.10.2015)

Browsset: NICE Guidance → Conditions and diseases → Mental health and behavioural conditions → anxiety

Fundne referencer: 5

NICE guidelines

- [Common mental health disorders_\(CG123\)_May 2011](#)
- [Generalised anxiety disorder and panic disorder \(with or without agoraphobia\) in adults_\(CG113\) January 2011](#)
- [Obsessive-compulsive disorder_\(CG31\)_November 2005](#)
- [Post-traumatic stress disorder \(PTSD\)_\(CG26\)_March 2005](#)
- [Social anxiety disorder: recognition, assessment and treatment_\(CG159\)_May 2013](#)

NICE technology appraisal guidance

- [Computerised cognitive behaviour therapy for depression and anxiety_\(TA97\)_February 2006](#)

NICE commissioning guides

- [Commissioning stepped care for people with common mental health disorders_\(CMG41\)_November 2011](#)

NICE quality standard

- [Anxiety disorders_\(QS53\)_February 2014](#)

NICE advice

- [First-choice antidepressant use in adults with depression or generalised anxiety disorder_\(KTT8\) January 2015](#)
- [Generalised anxiety disorder: quetiapine_\(ESUOM12\)_May 2013](#)

in development

- [Generalised anxiety disorder - quetiapine \[ID347\]_\(GID-TAG421\)_TBC Technology appraisals](#)

National Guideline Clearinghouse (USA) (IE)

(søgt d. 19.10.2015)

Browsset: [Disease/Condition](#) > [Psychiatry and Psychology](#) (405) > [Mental Disorders](#) (278) > [Anxiety Disorders](#) (23)

Fundne referencer: 23

1. [Screening, assessment, and care of anxiety and depressive symptoms in adults with cancer: an American Society of Clinical Oncology guideline adaptation.](#) 2014 May 20. Compare
 NGC:010560 Selected
 American Society of Clinical Oncology - Medical Specialty Society. [View all guidelines by the developer\(s\)](#)
2. [Social anxiety disorder: recognition, assessment and treatment.](#) 2013 May. Compare

- | | |
|--|--------------------------------------|
| NGC:009892 | Selected |
| National Collaborating Centre for Mental Health - National Government Agency [Non-U.S.]. View all guidelines by the developer(s) | <input type="checkbox"/> |
| 3. Practice parameter on gay, lesbian or bisexual sexual orientation, gender nonconformity, and gender discordance in children and adolescents. 2012 Sep. | Compare |
| NGC:009316 | Selected |
| American Academy of Child and Adolescent Psychiatry - Medical Specialty Society. View all guidelines by the developer(s) | <input type="checkbox"/> |
| 4. Management of perinatal mood disorders. A national clinical guideline. 2002 Jun (revised 2012 Mar). NGC:009026 | Compare |
| Scottish Intercollegiate Guidelines Network - National Government Agency [Non-U.S.]. View all guidelines by the developer(s) | Selected
<input type="checkbox"/> |
| 5. Common mental health disorders. Identification and pathways to care. 2011 May. | Compare |
| NGC:008755 | Selected |
| National Collaborating Centre for Mental Health - National Government Agency [Non-U.S.]. View all guidelines by the developer(s) | <input type="checkbox"/> |
| 6. Generalised anxiety disorder and panic disorder (with or without agoraphobia) in adults. Management in primary, secondary and community care. 2004 Dec (revised 2011 Jan). NGC:008660 | Compare |
| National Collaborating Centre for Mental Health - National Government Agency [Non-U.S.]; National Collaborating Centre for Primary Care - National Government Agency [Non-U.S.]. View all guidelines by the developer(s) | Selected
<input type="checkbox"/> |
| 7. Putting evidence into practice: evidence-based interventions for anxiety. 2008 Oct 1 (reaffirmed 2011). NGC:007728 | Compare |
| Oncology Nursing Society - Professional Association. View all guidelines by the developer(s) | Selected
<input type="checkbox"/> |
| 8. Use of psychiatric medications during pregnancy and lactation. 2008 Apr (reaffirmed 2012). NGC:006421 | Compare |
| American College of Obstetricians and Gynecologists - Medical Specialty Society. View | Selected
<input type="checkbox"/> |

all guidelines by the developer(s)

- | | | |
|-----|--|---|
| 9. | <p><u>Evidence-based guideline: assessment and management of psychiatric disorders in individuals with MS. Report of the Guideline Development Subcommittee of the American Academy of Neurology.</u> 2014 Jan 14. NGC:010200</p> <p>American Academy of Neurology - Medical Specialty Society. <i>View all guidelines by the developer(s)</i></p> | Compare
Selected
<input type="checkbox"/> |
| 10. | <p><u>Adult depression in primary care.</u> 1996 Jan (revised 2013 Sep). NGC:010042</p> <p>Institute for Clinical Systems Improvement - Nonprofit Organization. <i>View all guidelines by the developer(s)</i></p> | Compare
Selected
<input type="checkbox"/> |
| 11. | <p><u>Best evidence statement (BEST). Oral anxiolytic medication prior to ambulatory healthcare encounters for individuals with special developmental and behavioral challenges.</u> 2011 Mar 10. NGC:008546</p> <p>Cincinnati Children's Hospital Medical Center - Hospital/Medical Center. <i>View all guidelines by the developer(s)</i></p> | Compare
Selected
<input type="checkbox"/> |
| 12. | <p><u>Best evidence statement (BEST). In children with a history of child abuse or neglect does preparation for medical procedures using medical play vs. no preparation reduce anxiety?</u> 2010 Oct 26. NGC:008172</p> <p>Cincinnati Children's Hospital Medical Center - Hospital/Medical Center. <i>View all guidelines by the developer(s)</i></p> | Compare
Selected
<input type="checkbox"/> |
| 13. | <p><u>Deep brain stimulation for obsessive-compulsive disorder: systematic review and evidence-based guideline sponsored by the American Society for Stereotactic and Functional Neurosurgery and the Congress of Neurological Surgeons (CNS) and endorsed by the CNS and American Association of Neurological Surgeons.</u> 2014 Oct. NGC:010550</p> <p>Congress of Neurological Surgeons - Professional Association. <i>View all guidelines by the developer(s)</i></p> | Compare
Selected
<input type="checkbox"/> |
| 14. | <p><u>Mental illness & stress.</u> 2004 (revised 2013 Nov 18). NGC:010123</p> <p>Work Loss Data Institute - For Profit Organization. <i>View all guidelines by the developer(s)</i></p> | Compare
Selected |

- | | | |
|-----|---|---|
| | | <input type="checkbox"/> |
| 15. | Clinical Pharmacogenetics Implementation Consortium guideline for <i>CYP2D6</i> and <i>CYP2C19</i> genotypes and dosing of tricyclic antidepressants. 2013 May. NGC:009742
Clinical Pharmacogenetics Implementation Consortium - Independent Expert Panel.
<i>View all guidelines by the developer(s)</i> | Compare
Selected
<input type="checkbox"/> |
| 16. | Best evidence statement (BEST). Psycho-social interventions for primary caregivers of newly diagnosed pediatric oncology patients. 2012 Nov 28. NGC:009492
Cincinnati Children's Hospital Medical Center - Hospital/Medical Center. <i>View all guidelines by the developer(s)</i> | Compare
Selected
<input type="checkbox"/> |
| 17. | Practice parameter for the assessment and treatment of children and adolescents with obsessive-compulsive disorder. 2012 Jan. NGC:008982
American Academy of Child and Adolescent Psychiatry - Medical Specialty Society.
<i>View all guidelines by the developer(s)</i> | Compare
Selected
<input type="checkbox"/> |
| 18. | VA/DoD clinical practice guideline for management of post-traumatic stress. 2004 Jan (revised 2010 Oct). NGC:008249
Department of Defense - Federal Government Agency [U.S.]; Department of Veterans Affairs - Federal Government Agency [U.S.]; Veterans Health Administration - Federal Government Agency [U.S.]. <i>View all guidelines by the developer(s)</i> | Compare
Selected
<input type="checkbox"/> |
| 19. | Practice guideline for the treatment of patients with obsessive-compulsive disorder. 2007 (reaffirmed 2012). NGC:005841
American Psychiatric Association - Medical Specialty Society. <i>View all guidelines by the developer(s)</i> | Compare
Selected
<input type="checkbox"/> |
| 20. | Brain injury rehabilitation in adults. A national clinical guideline. 2013 Mar. NGC:009702
Scottish Intercollegiate Guidelines Network - National Government Agency [Non-U.S.]. <i>View all guidelines by the developer(s)</i> | |
| 21. | Practice parameter for psychodynamic psychotherapy with children. 2012 May. NGC:009070
American Academy of Child and Adolescent Psychiatry - Medical Specialty | Compare
Selected
<input type="checkbox"/> |

Society. [View all guidelines by the developer\(s\)](#)

22. [Eating disorders during pregnancy and postpartum](#). 2007 May 3 (revised 2010 Sep 16).

NGC:008251

University of Arkansas for Medical Sciences, ANGELS (Antenatal & Neonatal Guidelines, Education and Learning System) - Academic Institution. [View all guidelines by the developer\(s\)](#)

Scottish Intercollegiate Guidelines Network (SIGN)

(søgt d. 19.10.2015)

Browsset: By topic → Mental Health

Fundne referencer: 0

HTA Databasen (CRD database)

(søgt d. 19.10.2015)

Results for: (anxiety disorder* OR anxious OR anxiety* OR social phobia OR social anxiety) IN HTA FROM 2005 TO 2015

Fundne referencer: 94 (37 Overført til Covidence)

Results for: ((anxiety*)) and (Full publication record:ZDT) IN HTA FROM 2005 TO 2015

First	1	2	Last	Show all previews	Select all	Clear selections	Export
	Year	Database	Source	Title			
<input type="checkbox"/>	2015	HTA	Health Quality Ontario (HQO)	Cognitive-behavioural therapy for anxiety and depression in patients with chronic obstructive pulmonary disease (COPD): a			

				rapid review [Preview]	
<input type="checkbox"/>	2015	HTA	Institute for Clinical Evaluative Sciences (ICES)	The mental health of children and youth in Ontario: a baseline scorecard [Preview]	
<input type="checkbox"/>	2015	HTA	NIHR Health Technology Assessment programme	Treatment of childhood anxiety disorder in the context of maternal anxiety disorder: a randomised controlled trial and economic analysis [Preview]	
<input type="checkbox"/>	2014	HTA	NIHR Health Technology Assessment programme	Clinical effectiveness of interventions for treatment-resistant anxiety in older people: a systematic review [Preview]	
<input type="checkbox"/>	2014	HTA	NIHR Public Health Research (PHR) programme	Crime, fear of crime and mental health: Synthesis of theory and systematic reviews of interventions and qualitative evidence [Preview]	
<input type="checkbox"/>	2014	HTA	Agency for Healthcare Research and Quality (AHRQ)	Meditation programs for psychological stress and well-being [Preview]	
<input type="checkbox"/>	2014	HTA	Canadian Agency for Drugs and Technologies in Health (CADTH)	Neurofeedback and biofeedback for mood and anxiety disorders: a review of the clinical evidence and guidelines [Preview]	
<input type="checkbox"/>	2014	HTA	Canadian Agency for Drugs and Technologies in Health (CADTH)	Neuro-linguistic programming for the treatment of adults with post-traumatic stress disorder, general anxiety disorder, or depression: a review of clinical effectiveness and guidelines [Preview]	

<input type="checkbox"/>	2014	HTA	NIHR Horizon Scanning Centre (NIHR HSC)	Pesky gNATs cognitive behaviour therapy for children with anxiety and low mood [Preview]	
<input type="checkbox"/>	2014	HTA	Canadian Agency for Drugs and Technologies in Health (CADTH)	Short- and long-term use of benzodiazepines in patients with generalized anxiety disorder: a review of guidelines [Preview]	
<input type="checkbox"/>	2014	HTA	Canadian Agency for Drugs and Technologies in Health (CADTH)	Transcranial magnetic stimulation for the treatment of adults with PTSD, GAD, or depression: a review of clinical effectiveness and guidelines [Preview]	
<input type="checkbox"/>	2013	HTA	The Swedish Council on Health Technology Assessment (SBU)	[Internet-based psychological treatments for anxiety and mood disorders] [Preview]	
<input type="checkbox"/>	2013	HTA	Institute for Clinical Effectiveness and Health Policy (IECS)	[Music therapy - update on the evidence about its indications] [Preview]	
<input type="checkbox"/>	2013	HTA	Health Quality Ontario (HQO)	Patient experiences of depression and anxiety with chronic disease: a systematic review and qualitative meta-synthesis [Preview]	
<input type="checkbox"/>	2013	HTA	Canadian Agency for Drugs and Technologies in Health (CADTH)	The emotional freedom technique for the treatment of post-traumatic stress disorder, depression, or anxiety: clinical evidence [Preview]	
<input type="checkbox"/>	2013	HTA	Canadian Agency for Drugs and Technologies in Health (CADTH)	Transcendental meditation for posttraumatic stress disorder, depression, and anxiety: a review of clinical effectiveness [Preview]	

<input type="checkbox"/>	2012	HTA	Canadian Agency for Drugs and Technologies in Health (CADTH)	Neurofeedback and biofeedback for post-traumatic stress disorder, generalized anxiety disorder, and depression: a review of the clinical evidence and guidelines. [Preview]	
<input type="checkbox"/>	2012	HTA	The Regional Health Technology Assessment Centre (HTA-centrum)	Psychological treatment of dental anxiety among adults [Preview]	
<input type="checkbox"/>	2011	HTA	National Horizon Scanning Centre (NHSC)	Agomelatine (Valdoxan) for generalised anxiety disorder - first or second line [Preview]	
<input type="checkbox"/>	2011	HTA	Norwegian Knowledge Centre for the Health Services (NOKC)	[Effect of vitamins, minerals and other dietary supplements on mental health symptoms for people with ADHD, anxiety disorders, bipolar disorder or depression] [Preview]	
<input type="checkbox"/>	2011	HTA	National Institute for Health and Clinical Excellence (NICE)	Generalised anxiety disorder and panic disorder (with or without agoraphobia) in adults [Preview]	
<input type="checkbox"/>	2011	HTA	Agency for Healthcare Research and Quality (AHRQ)	Off-label use of atypical antipsychotics: an update [Preview]	
<input type="checkbox"/>	2011	HTA	Norwegian Knowledge Centre for the Health Services (NOKC)	[The effects of group education on patients and their next of kin] [Preview]	
<input type="checkbox"/>	2009	HTA	Canadian Agency for Drugs and Technologies in Health (CADTH)	Group therapy for anxiety: a review of the clinical effectiveness [Preview]	
<input type="checkbox"/>	2008	HTA	National Horizon Scanning Centre (NHSC)	Saregutant (SR-48968) for depression and anxiety [Preview]	
<input type="checkbox"/>	2007	HTA	The Swedish Council on Health Technology Assessment (SBU)	Computer-based cognitive behavioral therapy for anxiety disorders or depression - early assessment briefs (Alert) [Preview]	

2007 HTA Agency for Healthcare Research and Quality (AHRQ) Meditation practices for health: state of the research [\[Preview\]](#)

2006 HTA NIHR Health Technology Assessment programme Amniocentesis results: investigation of anxiety. The ARIA trial. [\[Preview\]](#)

2006 HTA National Institute for Health and Clinical Excellence (NICE) Computerised cognitive behaviour therapy for depression and anxiety: review of Technology Appraisal 51 [\[Preview\]](#)

2006 HTA NIHR Health Technology Assessment programme Computerised cognitive behaviour therapy for depression and anxiety update: a systematic review and economic evaluation [\[Preview\]](#)

2006 HTA New Zealand Health Technology Assessment (NZHTA) Effective models of mental health service provision and workforce configuration in the primary care setting [\[Preview\]](#)

2006 HTA Danish Centre for Evaluation and Health Technology Assessment (DACEHTA) [Preventive outpatient treatment in affective disorders. Results from a health technology assessment] [\[Preview\]](#)

2006 HTA Institute for Clinical Effectiveness and Health Policy (IECS) Usefulness of music therapy in clinical practice [\[Preview\]](#)

2005 HTA NIHR Health Technology Assessment programme A trial of problem-solving by community mental health nurses for anxiety, depression and life difficulties among general practice patients. The CPN-GP study [\[Preview\]](#)

2005 HTA NIHR Health Technology Assessment programme Long-term outcome of cognitive behaviour therapy clinical trials in central Scotland [\[Preview\]](#)

2005 HTA New Zealand Health Technology Assessment (NZHTA) The effectiveness of mental health promotion, prevention and early intervention in children, adolescents and adults: a critical appraisal of the literature [\[Preview\]](#)

<input type="checkbox"/>	2005 HTA	The Swedish Council on Health Technology Assessment (SBU)	Treatment of anxiety disorders: a systematic

SBU, Sverige (IE)

(søgt 19-10-2015)

(SBU utvärdering, SBU övriga publikationer, Regional utvärdering; 2005-2015)

Ångeststörning, ångest, social fobi, social ångest, generaliserat ångestsyndrom, Anxiety, anxiety disorder, social phobia, social anxiety, generalised anxiety disorder

Fundne referencer: 5

Internetförmedlad psykologisk behandling vid ångest- och förstämningssyndrom

Sammanfattning och slutsatser

Psykologisk behandling rangordnas högt i nationella riktlinjer för behandling av flera ångest- och förstämningssyndrom, och utgör en central del av behandlingsutbudet vid dessa tillstånd. Tillgängligheten till behandling varierar dock kraftigt mellan landstingen.

Traditionellt har psykologisk behandling förmedlats individuellt eller i grupp av en terapeut. Internet har inneburit nya möjligheter för att förmedla behandling på distans, i syfte att öka tillgängligheten och nå grupper som av olika skäl inte får behandling inom hälso- och sjukvården. I synnerhet internetförmedlad kognitiv beteendeterapi (KBT) har utvärderats i ett stort antal studier under drygt ett decennium.

Internetförmedlad behandling är vanligtvis strukturerad som ett självhjälpssystem under omkring 10 veckors tid med stöd av en kvalificerad behandlare via mejl, via behandlingsplattformen eller per telefon. Vissa program utförs dock helt självständigt eller med administrativt stöd. Deltagaren får via en webbsida tillgång till ett textavsnitt i veckan och övningar att genomföra på egen hand. Programmen varierar avseende interaktivitet och förekomst av ljud- och videoklipp.

SBU har utvärderat nytta och risk av internetförmedlad psykologisk behandling på kort och lång sikt vid ångest- och förstämningssyndrom för barn, ungdomar och vuxna. Utvärderingen inbegriper också behandlingsformens kostnadseffektivitet samt aspekter av etik, överförbarhet och implementering baserat på den vetenskapliga litteraturen.

Program för att förebygga psykisk ohälsa hos barn

Denna rapport publicerades 2010. Nya studier kan ha tillkommit. [Läs mer](#) »

Slutsatser

- Av 33 bedömda standardiserade och strukturerade insatser (program) som syftar till att förebygga psykisk ohälsa hos barn har sju ett begränsat vetenskapligt stöd i den internationella litteraturen. Det är föräldrastödsprogrammen Incredible Years, Triple-P, familjestödsprogrammet Family Check Up samt skolprogrammen Good Behavior Game, Coping Power, Coping With Stress och FRIENDS. Effekterna är med få undantag små. Studierna är utförda i andra länder.

Eftersom effekterna sannolikt varierar med sociala och kulturella sammanhang är det oklart i vilken utsträckning som programmen kan överföras till Sverige med bibehållen effekt. Programmen kan också behöva anpassas så att de överensstämmer med svenska värderingar och syn på barns rätt.

- I Sverige används ett hundratal olika program för att förebygga psykisk ohälsa hos barn, i huvudsak av utagerande typ. Inget av dem har utvärderats i Sverige i randomiserade studier med minst sex månaders uppföljning.

Programmen De Otroliga Åren (översatt från Incredible Years), Triple-P och Family Check Up har enligt internationella studier begränsat vetenskapligt stöd för förebyggande effekt.

Programmen KOMET, COPE, SET, StegVis, Beardslee's familjeintervention, Connect och DISA har undersökts i minst en kontrollerad studie vardera men har inte tillräckligt vetenskapligt stöd för förebyggande effekt.

Övriga program som används i Sverige är inte vetenskapligt utprovade som preventionsprogram.

- Program som bygger på att ungdomar med utagerande problem träffas i grupp kan öka risken för normbrytande beteenden. Andra negativa effekter för såväl program för utagerande som för inåtvända problem är tänkbara men ofullständigt belysta.
- Det behövs randomiserade studier som undersöker om de program som används har förebyggande effekt i svenska populationer och inte medför risker. Det behövs också hälsoekonomiska studier som undersöker om programmen är kostnadseffektiva.



[Till rapporten i fulltext \(pdf, 2MB\) »](#)

[Rapporten uppdelad per kapitel »](#)

Datorbaserad kognitiv beteendeterapi vid ångestsyndrom eller depression

En uppdatering av denna rapport publicerades 2013. [Läs rapporten »](#)

Sammanfattning och slutsatser

Metod och målgrupp

Ångest och depression är två tillstånd som drabbar en stor del av befolkningen. Ångest innebär en intensiv ångslan och känsla av olust kopplad till en förväntad fara eller olycka. Utmärkande för symtomen är att de är av både fysisk och psykisk karaktär. Med ångestsyndrom avses att flera symtom på ångest förekommer

samtidigt på ett specifikt sätt och med en viss varaktighet. Syndromen kan delas in i paniksyndrom, tvångssyndrom, posttraumatiskt stressyndrom, generaliserat ångestsyndrom, social fobi, specifika fobier samt ångestsyndrom UNS (utan närmare specifikation). Typiskt för depression är en känsla av hopplöshet, meningslöshet och nedstämdhet över en längre period. Liksom vid ångestsyndrom är patienter med depression ofta drabbade av andra tillstånd eller missbruk och en del blir oförmögna till arbete under kortare eller längre perioder.

Kognitiv beteendeterapi (KBT) är en form av psykologisk behandling som fokuserar på patientens tankar, känslor och beteende från ett inlärningsperspektiv, vilken har visat sig ha effekt vid ångestsyndrom respektive depression. Inom KBT betonas betydelsen av självhjälp och utvecklande av förmågan att lära sig handskas med sina problem. Datorbaserad KBT innebär att patienten går igenom en behandling som är baserad på KBT-manualer och som har anpassats till ett datorformat. I de flesta fall innefattar den datorbaserade behandlingen någon form av personlig kontakt med en KBT-terapeut, som följer patientens framsteg och finns tillgänglig för att svara på frågor, exempelvis via e-post, telefonsamtal eller vid få, korta personliga möten. I vissa av de datorbaserade programmen är kontakten med en KBT-terapeut minimal, medan andra mer används som ett komplement till KBT som ges av en terapeut. Det är inte klarlagt vilka kriterier som bör tillämpas för att välja ut patienter med depression eller ångestsyndrom som kan vara lämpliga för behandling med datorbaserad KBT. I dagsläget går det således inte att uppskatta storleken på den potentiella målgruppen för metoden.

Behandling av ångestsyndrom

Denna rapport publicerades 2005. Nya studier kan ha tillkommit. [Läs mer](#) »

SBU:s slutsatser

- För samtliga ångestsyndrom finns behandlingsmetoder med dokumenterad effekt (Evidensstyrka 1). Med undantag för specifika fobier är effekterna av såväl farmakologisk som psykoterapeutisk behandling måttliga. Symtomen lindras men det är sällan som full symtomfrihet uppnås. Med få undantag återkommer symtomen när behandlingen avslutats.
- Samhällskostnaderna, främst i form av nedsatt produktivitet, ökad sjuklighet och dödlighet och ökad konsumtion av somatisk vård är höga. Kostnadseffektiviteten för olika behandlingsalternativ är inte klarlagd.
- Det vetenskapliga underlaget för att jämföra såväl effekter som kostnadseffektivitet mellan olika behandlingsmetoder är otillräckligt.
- Studier av psykodynamiskt inriktade behandlingar saknas nästan helt.
- Några bensodiazepiner har dokumenterad effekt vid vissa ångestsyndrom. Samtidigt är det väl belagt att preparaten ger betydande problem i form av biverkningar, beroende eller att symtomen förvärras igen efter en tids behandling.

- Det finns inga studier som entydigt belyser varför ångestsyndrom är förknippat med överdödlighet. Långtidsstudier som undersöker huruvida man kan minska överdödligheten med hjälp av någon behandling saknas.

Utvärdering av effekten av mindfulness-baserade terapier vid stress, depression och ångest (2010)
Landsting/Region: Örebro läns landsting
Publicerad: 2010



Till rapporten »

Du kommer att lämna SBU:s webbplats

Socialstyrelsen, Sverige (IE)

(søgt d. 19.10.2015)

Browsed listen over Nationella riktlinjer for guidelines med ordene: ångest, ångeststörning, , social fobi, social ångest, generaliserat ångestsyndrom
Fundne referencer: 1

Nationella riktlinjer för depression och ångestsyndrom innehåller rekommendationer om effektivt omhändertagande, diagnostik, uppföljning och olika behandlingar som KBT, antidepressiva läkemedel och ECT.

<http://www.socialstyrelsen.se/nationellariktlinjerfordepressionochangest>

Helsedirektoratet, Norge (IE)

(søgt d. 19.10.2015)

Søgt på: angst, generalisert angstlidelse, angstlidelse, sosial fobi, sosial angst

Fundne referencer: 2

Prioriteringsveileder - psykisk helsevern for barn og unge (gjelder fra 1. november 2015)

Angstlidelser

[https://helsedirektoratet.no/retningslinjer/psykisk-helsevern-for-barn-og-unge/seksjon?Tittel=angstlidelser-](https://helsedirektoratet.no/retningslinjer/psykisk-helsevern-for-barn-og-unge/seksjon?Tittel=angstlidelser-9621#Angstlidelser - lettere (ikke rett))

9621#Angstlidelser - lettere (ikke rett)

<https://helsedirektoratet.no/retningslinjer/psykisk-helsevern-for-barn-og-unge/seksjon?Tittel=angstlidelser-9621>

Prioriteringsveileder - psykisk helsevern for voksne (gjelder fra 1. november 2015)

Angst og tvang

<https://helsedirektoratet.no/retningslinjer/psykisk-helsevern-for-voksne/seksjon?Tittel=angst-og-tvang-9640>

Kunnskapssenteret, Norge (IE)

(søgt d. 15.10.2015)

Browset: type: Rapport fra Kunnskapssenteret, → Tema: Barn & unge, Psykisk helse → År: 2005-2015 → dokumenter med et af følgende ord i titlen: angst

Fundne referencer: 4

Rapport fra Kunnskapssenteret - Systematisk oversikt

Effekten av vitaminer, mineraler og andre kosttilskudd på psykiske symptomer hos personer med ADHD, angstlidelser, bipolar lidelse eller depresjon

[ForsidenPublikasjoner](#)

Publisert 10.02.2011 Endret 27.11.2014

Berg RC, Smedslund G. Rapport fra Kunnskapssenteret nr. 03 – 2011. ISBN 978-82-8121-394-4 ISSN 1890-1298.

- Hovedfunn
- [Sammendrag](#)
- [Hele publikasjonen](#)

Mentale lidelser er svært utbredt og har store konsekvenser for de som rammes, både pga plagsomme symptomer og ofte pga nedsatt sosial og yrkesmessig fungering. De vanligste behandlingsmetodene er ulike former for medikamenter (psykofarmaka) og samtaleterapi. Kosttilskudd, som flerumettede fettsyrer og vitaminer, benyttes noen ganger, men effekten av slik behandling er omdiskutert.

I den foreliggende oversikten har vi forsøkt å besvare: Hva er effekten av vitaminer, mineraler og andre kosttilskudd på psykiske symptomer hos personer med ADHD (attention deficit hyperactivity disorder), angst-lidelser, bipolar lidelse eller depresjon?

Vi utarbeidet en kunnskapsoversikt over nyere systematiske oversikter. Vi inkluderte 11 oversikter, tre av høy metodisk kvalitet og åtte av moderat metodisk kvalitet. Intervensjonene i de inkluderte oversiktene var flerumettede fettsyrer, inositol, folat og vitamin B-6.

Rapport fra Kunnskapscenteret - Systematisk oversikt

Effekter av skolehelsetjenesten på barn og unges helse og oppvekstvilkår

[ForsidenPublikasjoner](#)

Publisert 30.06.2010 Endret 27.11.2014

Kristin Thuve Dahm, Brynjar Landmark, Ingvild Kirkehei og Liv Merete Reinar.. Rapport fra Kunnskapscenteret nr. 15 – 2010. ISBN 978-82-8121-355-5 ISSN 1890-1298.

- Hovedfunn
- [Sammendrag](#)
-

Bakgrunn: Helsestasjons- og skolehelsetjeneste er en lovpålagt tjeneste i Norge. Kommunene har ansvar og frihet til å organisere virksomheten på en hensiktsmessig måte, tilpasset lokale behov, så lenge gjeldende lover og forskrifter følges.

Oppdrag: Nasjonalt kunnskapscenter for helsetjenesten fikk i oppdrag fra Helsedirektoratet å oppsummere tilgjengelig forskning om effekten av skolehelsetjenesten på barn og unges helse og oppvekstvilkår. Denne kunnskapsoppsummeringen skal brukes som grunnlag for å utrede tiltak for å styrke skolehelsetjenesten.

Oppsummering av resultatene:

- Påvisning og tidlig henvisning til psykologisk behandling av barn med depresjon reduserer muligens depresjon hos barn og unge.
- Angstforebyggende tiltak bygget på kognitiv adferdsterapi har muligens en liten effekt på å redusere forekomst av angst hos skolebarn generelt og på å redusere angst hos skolebarn som allerede har symptomer på angst.
- Kombinasjon av undervisning og utdeling av prevensjon reduserer muligens antall uønskede svangerskap.
- Forebyggende røyketiltak i skolen synes å ha effekt på intensjon om å røyke, på å forebygge røykestart og på røykeadferd
- Det kan se ut til at skolebaserte programmer for barn med astma fører til økt kunnskap og mestring.

Oversikten kan bare delvis besvare vår problemstilling. Vi har ikke funnet noen systematiske oversikter som omhandler helsepersonell i en skolehelsetjeneste slik den er organisert i Norge, men vi har identifisert flere forebyggende og helsefremmende tiltak gjennomført på skolen hvor helsepersonell er involvert. Vi har inkludert tiltak innen psykisk helse, adferd og fysisk helse.

Rapport fra Kunnskapscenteret - Systematisk oversikt

Ambulante psykiatriske helsetjenester til barn og ungdom som alternativ til institusjonsbehandling

[ForsidenPublikasjoner](#)

Publisert 03.10.2014 Endret 28.11.2014

Rapport fra Kunnskapscenteret nr. 22 – 2006. ISBN 82-8121-129-6 ISSN 1890-1298.

- Sammendrag
- [Hele publikasjonen](#)

Behandling av psykiske vansker og atferdsproblemer hos barn og unge ses i sammenheng med pasientens nærmiljø hvor familien, skolen og venner har stor betydning i forhold til sykdomsutvikling og

behandling. Dette har ført til utvikling av flere nærmiljøbaserte metoder og organisatoriske modeller for behandling av denne pasientgruppen. Ambulant psykiatrisk behandling er tatt i bruk i Norge, og det er politisk vilje til å støtte videre utbygging av ambulante tjenester også for barn og unge. Dette er i tråd med internasjonal utvikling innen barne- og ungdomspsykiatrien.

Denne rapporten er bestilt fra Regionsenter for barn og unges psykiske helse, Helseregion øst og sør (R.BUP), som del av en større utredning om bruk av nærmiljøbaserte tjenester i behandling av barn og ungdom med alvorlige psykiske problemer. Rapporten omhandler forskningsbasert kunnskap om ambulant behandling av barn og ungdom i alderen 4-18 år med alvorlige psykiske problemer og atferdsvansker. Rapporten gir en oversikt over metoder og modeller for ambulante psykiatriske tjenester som har vært evaluert gjennom kontrollerte studier.

Vi søkte etter systematiske oversikter, randomiserte kontrollerte studier, kontrollerte før- og etter-studier og tidsserier i internasjonale databaser i juni og juli 2006. Studier som oppfylte de forhåndsbestemte kriteriene ble kritisk vurdert og sammenfattet.

Rapport fra Kunnskapscenteret - Oversikt over systematiske oversikter

Effekt av psykososiale tiltak for forebygging og behandling av depresjon og angst blant risikoutsatte barn og ungdommer

[ForsidenPublikasjoner](#)

Publisert 24.11.2014 Endret 26.11.2014

Munthe-Kaas HM, Johansen S, Blaasvær N, Hammerstrøm KT, Nilsen W. Rapport fra Kunnskapscenteret nr. 22 – 2014. ISBN 978-82-8121-846-8 ISSN 1890-1298 .

- Hovedfunn
- [Sammendrag](#)
- [Hele publikasjonen](#)

Relativt mange barn og unge har milde symptomer på depresjon eller angst. Når symptomene påvirker livskvaliteten kan de trenge profesjonell hjelp for å bli friske. Psykososiale tiltak for depresjon og angst er ikke-medikamentelle tiltak som kan rettes mot barn fra risikogrupper (forebyggende tiltak), eller de med tydelig symptomer på depresjon eller angst og dem som er fått en diagnose på depresjon eller angst (behandlingstiltak).

Netpunkt

Søgt d. 19.10.2015

(angst? Eller social fobi Eller social angst?) Og (retningslin? Eller guideline? Eller anbefal? Eller referenceprogram?)

2 fund

Faglige retningslinjer for henvisning til psykolog. : For pasienter med let til moderat depression eller let til moderat angst

Emne: [faglige retningslinjer](#) ; [depression](#) ; [angst](#) ; [almen praksis](#) ; [praktiserende psykolog](#) ; [Faglig rådgivning](#)

BOG: Version: 1.0; Versionsdato: 23. februar 2012. - Kbh., 2012. - 22 s.

Referenceprogram for angstlidelser hos voksne : udarbejdet af en arbejdsgruppe nedsat af Sekretariatet for Referenceprogrammer – SfR

Emne: [Referenceprogram](#) ; [kliniske retningslinjer](#) ; [psykiatriområdet](#) ; [Angstlidelser](#) ; [angst](#) ; [Anbefaling](#)

BOG: Version: 1.0. - Kbh., Sundhedsstyrelsen, 2007. - 158 sider

Medline

(Søgt d. 28.10.2015)

Database(s): **Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations and Ovid MEDLINE(R)** 1946 to Present

Search Strategy:

#	Searches	Results
1	Anxiety Disorders/	25261
2	anxiety disorder*.mp.	36736
3	anxiety.mp.	167996
4	anxious.mp.	12328
5	social phob*.mp.	3486
6	social anxiety.mp.	3849
7	generalized anxiety disorder*.mp.	4313
8	generalised anxiety disorder*.mp.	436
9	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8	172586
10	(child* or infant* or adoles* or teenager* or teen* or youth or young* or paediatr* or pediatr* or school or juvenil*).mp.	3979444
11	9 and 10	59974
12	limit 11 to (yr="2005 -Current" and (guideline or practice guideline))	38
13	clinical guideline*.mp.	8115
14	exp Practice Guideline/ or exp Guideline/ or practice guideline.mp.	29702
15	13 or 14	37159
16	11 and 15	109
17	12 or 16	109
18	limit 17 to (yr="2005 -Current" and (danish or english or norwegian or swedish))	72

Embase

(Søgt d. 28.10.2015)

Database(s): **Embase** 1974 to 2015 October 27

Search Strategy:

#	Searches	Results
1	anxiety disorder/ or anxiety disorder*.mp.	64300
2	exp social phobia/ or social phobia.mp.	9868
3	generalised anxiety disorders.mp. or exp generalized anxiety disorder/	6885
4	social anxiety.mp.	5242
5	(generalized anxiety disorder* or generalised anxiety disorder*).mp.	9745
6	social phob*.mp.	9996
7	anxiety.ti,ab.	177311
8	anxious.ti,ab.	16980
9	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8	206138
10	(child* or infant* or adoles* or teenager* or teen* or youth or young* or paediatr* or peditr* or school or juvenil*).mp.	3898359
11	9 and 10	57544
12	practice guideline/	265060
13	clinical guideline*.mp.	11766
14	12 or 13	270817
15	11 and 14	635
16	limit 15 to ((danish or english or norwegian or swedish) and yr="2005 -Current")	475

PsycInfo

(Søgt d. 28.10.2015)

Database(s): **PsycINFO** 1806 to October Week 3 2015

Search Strategy:

#	Searches	Results
1	Anxiety Disorders/ or anxiety disorder*.mp.	31947
2	social phobia.mp. or exp Social Phobia/	6442
3	(generalized anxiety disorder* or generalised anxiety disorder*).mp.	6409
4	exp Generalized Anxiety Disorder/	2069
5	(social anxiety or social phob*).mp.	11477

6	anxious.ab,ti.	16837
7	anxiety.ab,ti.	146909
8	1 or 2 or 3 or 4 or 5 or 6 or 7	157754
9	(child* or infant* or adoles* or teenager* or teen* or youth or young* or paediatr* or pediatr* or school or juvenil*).mp.	1074845
10	8 and 9	45840
11	exp Treatment Guidelines/	5013
12	clinical guideline*.mp.	1655
13	practice guideline*.mp.	3052
14	11 or 12 or 13	8500
15	10 and 14	123
16	limit 15 to ((danish or english or norwegian or swedish) and yr="2005 -Current")	93

CINAHL

(Søgt d. 28.10.15)

#	Query	Limiters/Expanders	Results
S14	S10 AND S13	Limiters - Published Date: 20050101- 20151231; Language: Danish, English, Norwegian, Swedish Search modes - Boolean/Phrase	63
S13	S11 OR S12	Search modes - Boolean/Phrase	41,210
S12	"clinical guideline*"	Search modes - Boolean/Phrase	2,576
S11	(MH "Practice Guidelines") OR "practice guideline*"	Search modes - Boolean/Phrase	39,762
S10	S8 AND S9	Search modes - Boolean/Phrase	10,381
S9	(child* or infant* or adoles* or teenager* or teen* or youth or young* or paediatr* or pediatr* or school or juvenil*)	Search modes - Boolean/Phrase	636,431
S8	S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7	Search modes - Boolean/Phrase	30,398
S7	TI anxious OR AB anxious	Search modes - Boolean/Phrase	2,210

S6	TI anxiety OR AB anxiety	Search modes - Boolean/Phrase	27,935
S5	"Generalised Anxiety Disorder*"	Search modes - Boolean/Phrase	107
S4	(MH "Generalized Anxiety Disorder") OR "generalized anxiety disorder*"	Search modes - Boolean/Phrase	649
S3	"social anxiety*"	Search modes - Boolean/Phrase	658
S2	(MH "Social Anxiety Disorders") OR "social phob*"	Search modes - Boolean/Phrase	629
S1	(MH "Anxiety Disorders") OR "anxiety disorder*"	Search modes - Boolean/Phrase	5,415

777 referencer er overført til RefWorks. Efter dubletfjernelse er der overført 684 referencer til Covidence. Herunder over er der 39 guidelines i dokumentet, som ikke er overført til Covidence.

Supplerende søgning på Separation Anxiety

(14-01-2016)

Database(s): **Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations and Ovid MEDLINE(R) 1946 to Present**

Search Strategy:

#	Searches	Results
1	Anxiety Disorders/	26571
2	anxiety disorder*.mp.	37796
3	anxiety.mp.	166972
4	anxious.mp.	12210
5	social phob*.mp.	3440
6	social anxiety.mp.	3779
7	generalized anxiety disorder*.mp.	4237
8	generalised anxiety disorder*.mp.	434
9	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8	171522
10	(child* or infant* or adoles* or teenager* or teen* or youth or young* or paediatr* or peditr* or school or juvenil*).mp.	3928695
11	9 and 10	59397

12	limit 11 to (yr="2005 -Current" and (guideline or practice guideline))	37
13	clinical guideline*.mp.	8121
14	exp Practice Guideline/ or exp Guideline/ or practice guideline.mp.	29482
15	13 or 14	36954
16	11 and 15	105
17	12 or 16	105
18	limit 17 to (yr="2005 -Current" and (danish or english or norwegian or swedish))	68
19	exp Anxiety, Separation/	1943
20	separation anxiet*.mp.	1198
21	19 or 20	2628
22	10 and 21	2107
23	limit 22 to (yr="2005 -Current" and (guideline or practice guideline))	1
24	15 and 22	1
25	23 or 24	1
26	18 or 25	68

Ingen nye referencer

Database(s): **Embase** 1996 to 2016 Week 02

Search Strategy:

#	Searches	Results
1	anxiety disorder/ or anxiety disorder*.mp.	60299
2	exp social phobia/ or social phobia.mp.	9435
3	generalised anxiety disorders.mp. or exp generalized anxiety disorder/	7019
4	social anxiety.mp.	5010
5	(generalized anxiety disorder* or generalised anxiety disorder*).mp.	9251
6	social phob*.mp.	9477
7	anxiety.ti,ab.	149999
8	anxious.ti,ab.	13529
9	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8	175510
10	(child* or infant* or adoles* or teenager* or teen* or youth or young* or paediatr* or peditr* or school or juvenil*).mp.	2494223
11	9 and 10	50818
12	practice guideline/	259870
13	clinical guideline*.mp.	11520

14	12 or 13	265472
15	11 and 14	646
16	limit 15 to ((danish or english or norwegian or swedish) and yr="2005 -Current")	493
17	exp separation anxiety/	2530
18	separation anxiet*.mp.	2837
19	17 or 18	2837
20	10 and 19	2240
21	14 and 20	26
22	limit 21 to ((danish or english or norwegian or swedish) and yr="2005 -Current")	21
23	22 not 16	3

3 nye referencer

Database(s): **PsycINFO** 2002 to January Week 1 2016

Search Strategy:

#	Searches	Results
1	Anxiety Disorders/ or anxiety disorder*.mp.	23149
2	social phobia.mp. or exp Social Phobia/	4655
3	(generalized anxiety disorder* or generalised anxiety disorder*).mp.	4880
4	exp Generalized Anxiety Disorder/	2101
5	(social anxiety or social phob*).mp.	8251
6	anxious.ab,ti.	9589
7	anxiety.ab,ti.	85866
8	1 or 2 or 3 or 4 or 5 or 6 or 7	92120
9	(child* or infant* or adoles* or teenager* or teen* or youth or young* or paediatr* or pediater* or school or juvenil*).mp.	560522
10	8 and 9	28896
11	exp Treatment Guidelines/	4821
12	clinical guideline*.mp.	1359
13	practice guideline*.mp.	2673
14	11 or 12 or 13	7678
15	10 and 14	114
16	limit 15 to ((danish or english or norwegian or swedish) and yr="2005 -Current")	95
17	exp Separation Anxiety/	563
18	separation anxiet*.mp.	1396

19	17 or 18	1396
20	9 and 19	1208
21	14 and 20	7
22	21 not 16	4
23	limit 22 to ((danish or english or norwegian or swedish) and yr="2005 -Current")	1

1 ny reference

Cinahl

(15-01-2016)

#	Query	Results
S20	s19 NOT s14	2
S19	S13 AND S18	3
S18	S9 AND S17	378
S17	S15 OR S16	464
S16	separation anxiet*	464
S15	(MH "Separation Anxiety")	349
S14	S10 AND S13	63
S13	S11 OR S12	41,379
S12	"clinical guideline*"	2,534
S11	(MH "Practice Guidelines") OR "practice guideline*"	39,968
S10	S8 AND S9	10,171
S9	(child* or infant* or adoles* or teenager* or teen* or youth or young* or paediatr* or pediater* or school or juvenil*)	636,320
S8	S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7	29,849
S7	TI anxious OR AB anxious	2,165
S6	TI anxiety OR AB anxiety	27,387
S5	"Generalised Anxiety Disorder*"	103
S4	(MH "Generalized Anxiety Disorder") OR "generalized anxiety disorder*"	624
S3	"social anxiety*"	650

S2	(MH "Social Anxiety Disorders") OR "social phob*"	625
S1	(MH "Anxiety Disorders") OR "anxiety disorder*"	5,329
2 nye referencer		

I alt er der fundet 6 nye referencer i den supplerende søgning.