

**This draft has not been adopted or endorsed by the European Commission. Any views expressed are the preliminary views of the Commission services and may not in any circumstances be regarded as stating an official position of the Commission. The information transmitted is intended only for the Member State or entity to which it is addressed for discussions and may contain confidential and/or privileged material**

**ANNEX**

Identification number of the feed additive	Name of the holder of authorisation	Additive	Composition, chemical formula, description, analytical method	Species or category of animal	Maximum age	Minimum dose	Maximum dose	Other provisions	End of period of authorisation
						chlamydo spores/kg body weight and day			
<b>Category: zootechnical additives. Functional group: other zootechnical additives (reduction of parasitic infective nematode larvae on pasture)</b>									
4d27	International Animal Health Products Pty Ltd, represented by GAB Consulting GmbH	<i>Duddingtonia flagrans</i> NCIMB 30336	<b>Additive composition</b> Preparation of <i>Duddingtonia flagrans</i> NCIMB 30336 containing a minimum of $5 \times 10^5$ chlamydo spores/g additive. Solid form.  <b>Characterisation of the active substance</b> Viable chlamydo spores of <i>Duddingtonia flagrans</i> NCIMB 30336  <b>Analytical method <sup>(1)</sup></b> For the identification of <i>Duddingtonia flagrans</i> NCIMB 30336: DNA based methods.	Dairy cows  Dairy cows of minor bovine species  Dairy sheep  Dairy goats	-	$3 \times 10^4$	-	1. In the directions for use of the additive and premixtures, the storage conditions and stability to heat treatment shall be indicated.  2. The minimum dose of $3 \times 10^4$ chlamydo spores/kg body weight and day equivalates to: - in standard domestic bovine animals of 650 kg body weight: $8.58 \times 10^5$ chlamydo spores/kg complete feedingstuff with a moisture content of 12% - in standard sheep and goats of 60 kg body weight: $1.32 \times 10^6$ chlamydo spores/kg	[10 years from the date of entry into force of this Regulation. To be completed by the Service responsible for the publication]

<sup>(1)</sup> Details of the analytical methods are available at the following address of the Reference Laboratory: [https://joint-research-centre.ec.europa.eu/eurl-fa-eurl-feed-additives/eurl-fa-authorisation/eurl-fa-evaluation-reports\\_en](https://joint-research-centre.ec.europa.eu/eurl-fa-eurl-feed-additives/eurl-fa-authorisation/eurl-fa-evaluation-reports_en)

			<p>For the enumeration of viable chlamydo spores of <i>Duddingtonia flagrans</i> NCIMB 30336 in the feed additive, premixtures and compound feed: the method using yeast mannitol agar (YMA) with streptomycin and chloramphenicol and a most probable number (MPN) for the enumeration.</p>					<p>complete feedingstuff with a moisture content of 12%</p> <p>3. The additive shall only be used in feed for grazing animals.</p> <p>4. The following shall be indicated on the label of the additive: ‘Recommended maximum dose: <math>3 \times 10^5</math> chlamydo spores/kg body weight and day’.</p> <p>5. For users of the additive and premixtures, feed business operators shall establish operational procedures and organisational measures to address potential risks resulting from their use. Where those risks cannot be eliminated by such procedures and measures, the additive and premixtures shall be used with personal breathing and skin protective equipment.</p>	
--	--	--	--	--	--	--	--	---	--