

ANNEX

Identification number of the additive	Name of the additive	Composition, chemical formula, description, analytical method.	Species or category of animal	Maximum age	Minimum content	Maximum content	Other provisions	End of period of authorisation
					mg active substance/kg of complete feed with a moisture content of 12%			
Category: Sensory additives. Functional group: Flavouring compounds								
2b255-eo	Laurel leaf essential oil	Additive composition Essential oil from the leaves of <i>Laurus nobilis</i> L. Liquid form Characterisation of the active substance Laurel leaf essential oil: Essential oil, as defined by the Council of Europe ¹ , obtained from leaves of <i>Laurus nobilis</i> L. by steam distillation and further condensation of volatile	Turkeys for fattening	-	-	10	1. The additive shall be incorporated into the feed in the form of a premixture. 2. In the directions for use of the additive and premixtures, the storage conditions and the stability to heat treatment	[10 years from the date of entry into force of this Regulation. To be completed by the OP]
			Chickens for fattening and minor poultry for fattening	-	-	8,5		
			Pigs for fattening	-	-	10		
			Pigs for fattening of minor <i>Suidae</i>	-	-	10		
			Calves for fattening	6 months	-	10		

¹ Natural sources of flavourings - Report No. 2 (2007).

		constituents and separation from the aqueous phase by decantation. CAS number: 8002-41-3 EINECS: 283-272-5 ² FEMA number: 2613 CoE number: 255 <u>Specifications</u> 1,8-Cineole (eucalyptol): 41-58% α-Terpinyl acetate: 5,5 - 12,5% Sabinene (4(10)-thujene): 4 - 11,5% α-Pinene (pin-2(3)-ene): 4,5 - 9% β-Pinene (pin-2(10)-ene): 2,8 - 6% Linalool: 2 - 5,5% Methyleugenol: maximum 4% Estragole: maximum 0,4% Elemicin: maximum 0,01% Analytical method³ For the determination of 1,8-cineole (phytochemical marker) in the feed additive: - gas chromatography coupled with flame ionisation detection (GC-FID)	Sheep and goats for fattening	-	-	10	shall be indicated. 3. Simultaneous use of laurel leaf essential oil with other feed additives is permitted provided that the combined content of <i>p</i> -allylalkoxybenzenes (methyleugenol, estragole, elemicin, safrole, myristicin, dillapiol, and apiole) in feed materials and compound feed remains below the content that would result from using the authorised feed additive with the	
			Cattle for fattening; other ruminants for fattening except sheep, goats for fattening and calves for fattening up to 6 months old; <i>Camelidae</i> for fattening	-	-	10		
			<i>Leporidae</i> for fattening	-	-	10		
			Salmonids and minor finfish except brood stock	-	-	10		
			Dogs	-	-	2		

² Here it only refers to the essential oil from the leaves of *Laurus nobilis* L.

³ Details of the analytical methods are available at the following address of the Reference Laboratory: https://joint-research-centre.ec.europa.eu/eurl-fa-eurl-feed-additives/eurl-fa-authorisation/eurl-fa-evaluation-reports_en.

			Cats	-	-	2	highest content of those substances at its maximum or recommended maximum content for the relevant animal species or category. 4. For users of the additive and premixtures, feed business operators shall establish operational procedures and organisational measures to address the potential risks resulting from their use. Where those risks cannot be eliminated by such procedures and measures, the additive and premixtures shall be used with	
--	--	--	------	---	---	---	--	--

							personal breathing, eye and skin protective equipment.	
--	--	--	--	--	--	--	--	--