

ANNEX I

Identifi-cation number of the additive	Name of the holder of authorisa-tion	Name of the additive	Composition, chemical formula, description, analytical method	Species or category of animal	Maxi-mum age	Mini-mum content	Maximu m content	Other provisions	End of period of authorisa-tion
						CFU/kg of complete feed with a moisture content of 12 %			
Category: zootechnical additives. Functional group: gut flora stabilisers.									
4b1900	Biochem Zusatzstoffe Handels- und Produktions ges. mbH	<i>Weizmannia faecalis</i> DSM 32016	<p>Additive composition Preparation of <i>Weizmannia faecalis</i> DSM 32016 containing a minimum of 2×10^{10} CFU/g additive Solid form</p> <p>Characterisation of the active substance Viable spores of cells of <i>Weizmannia faecalis</i> DSM 32016</p> <p>Analytical method (1) Identification: DNA sequencing methods or Pulsed Field Gel Electrophoresis (PFGE) – CEN/TS 17697. Enumeration in the feed additive, premixtures, compound feed and water for drinking: Spread plate method on MRS agar (based on EN</p>	<p>Suckling and weaned Suidae piglets</p> <p>Poultry for fattening</p> <p>Ornamental birds</p>	-	1×10^9	-	<p>1. In the directions for use of the additive and premixtures, the storage conditions and stability to heat treatment shall be indicated.</p> <p>2. The additive may be used simultaneously with the following coccidiostats, in accordance with their respective conditions of authorisation as feed additives: halofuginone, diclazuril, monensin sodium, robenidine hydrochloride, salinomycin sodium, amprolium, a combination of monensin sodium with narasin and a combination of narasin and nicarbazin.</p> <p>3. For users of the additive and premixtures, feed business operators shall establish operational procedures and organisational measures to address potential risks resulting from their use. Where those risks cannot be eliminated by such procedures and measures, the additive and premixtures shall be used with personal breathing and skin protective equipment.</p>	15.12.2030

(1) Details of the analytical methods are available at the following address of the Reference Laboratory: https://joint-research-centre.ec.europa.eu/eurl-fa-eurl-feed-additives/eurl-fa-authorisation/eurl-fa-evaluation-reports_en

		15787 method).							
--	--	----------------	--	--	--	--	--	--	--

ANNEX II

PART I

Identifi-cation number of the additive	Name of the holder of authorisa-tion	Name of the additive	Composition, chemical formula, description, analytical method	Species or category of animal	Maxi-mum age	Mini-mum content	Maxim um content	Mini-mum content	Maxim um content	Other provisions	End of perio d of autho-risa-tion
						CFU/kg of complete feed with a moisture content of 12 %	CFU/l of water for drinking				
Category: zootechnical additives. Functional group: gut flora stabilisers.											
4b1900	Biochem Zusatzstoffe Handels- und Produktions ges. mbH	<i>Weizmannia faecalis</i> DSM 32016	<p>Additive composition Preparation of <i>Weizmannia faecalis</i> DSM 32016 containing a minimum of 2×10^{10} CFU/g additive Solid form</p> <p>Characterisation of the active substance Viable spores of <i>Weizmannia faecalis</i> DSM 32016</p> <p>Analytical method (2) Identification: DNA sequencing methods or Pulsed Field Gel Electrophoresis (PFGE) – CEN/TS 17697. Enumeration in the feed additive,</p>	All poultry species reared for laying All poultry species reared for breeding	-	1×10^9	-	5×10^8	-	1. In the directions for use of the additive and premixtures, the storage conditions, the stability to heat treatment and the stability in water for drinking shall be indicated. 2. The additive may be used via water for drinking. 3. The additive may be used simultaneously with the following coccidiostats, in accordance with their respective conditions of authorisation as feed additives: halofuginone, diclazuril, monensin sodium, robenidine hydrochloride, salinomycin sodium, amprolium, a combination of monensin sodium with nicarbazin and a combination of narasin and	1 May 2034

(2) Details of the analytical methods are available at the following address of the Reference Laboratory: https://joint-research-centre.ec.europa.eu/eurl-fa-eurl-feed-additives/eurl-fa-authorisation/eurl-fa-evaluation-reports_en

			premixtures, compound feed and water for drinking: Spread plate method on MRS agar (based on EN 15787 method).						nicarbazin. 4. For users of the additive and premixtures, feed business operators shall establish operational procedures and organisational measures to address potential risks resulting from their use. Where those risks cannot be eliminated by such procedures and measures, the additive and premixtures shall be used with personal breathing and skin protective equipment.	
--	--	--	--	--	--	--	--	--	--	--

PART II

Identifi-cation number of the additive	Name of the holder of authorisa-tion	Name of the additive	Composition, chemical formula, description, analytical method	Species or category of animal	Maxi-mum age	Minimum content	Maximum content	Other provisions	End of perio-d of autho-risa-tion	
						CFU/l of water for drinking				
Category: zootechnical additives. Functional group: gut flora stabilisers.										
4b1900	Biochem Zusatzstoffe Handels- und Produktions ges. mbH	<i>Weizmannia faecalis</i> DSM 32016	<p>Additive composition Preparation of <i>Weizmannia faecalis</i> DSM 32016 containing a minimum of 2×10^{10} CFU/g additive Solid form</p> <p>Characterisation of the active substance Viable spores of <i>Weizmannia faecalis</i> DSM 32016</p>	All poultry species for fattening Suckling and weaned Suidae piglets Ornamental birds	-	5×10^8	-	1. In the directions for use of the additive, the storage conditions and the stability in water for drinking shall be indicated. 2. The additive shall be used via water for drinking. 3. The additive may be used simultaneously with the following coccidiostats, in accordance with their respective conditions of authorisation as feed additives: halofuginone, diclazuril,	1 May 2034	

		<p><i>Analytical method⁽³⁾</i></p> <p>Identification: DNA sequencing methods or Pulsed Field Gel Electrophoresis (PFGE) – CEN/TS 17697.</p> <p>Enumeration in the feed additive, premixtures, compound feed and water for drinking: Spread plate method on MRS agar (based on EN 15787 method).</p>					monensin sodium, robenidine hydrochloride, salinomycin sodium, amprolium, a combination of monensin sodium with nicarbazin and a combination of narasin and nicarbazin.	4. For users of the additive and premixtures, feed business operators shall establish operational procedures and organisational measures to address potential risks resulting from their use. Where those risks cannot be eliminated by such procedures and measures, the additive and premixtures shall be used with personal breathing and skin protective equipment.	
--	--	---	--	--	--	--	---	---	--

⁽³⁾ Details of the analytical methods are available at the following address of the Reference Laboratory: https://joint-research-centre.ec.europa.eu/eurl-fa-eurl-feed-additives/eurl-fa-authorisation/eurl-fa-evaluation-reports_en